


Scorecard for Accountability and Action



Malaria is present throughout Burkina Faso, with transmission most intense in the southern part of the country. The annual reported number of malaria cases in 2018 was 11,970,321 with 4,294 deaths.

Metrics

Commodities Financed and Financial Control		
LLIN financing 2020 projection (% of need)		97
Public sector RDT financing 2020 projection (% of need)		100
Public sector ACT financing 2020 projection (% of need)		93
World Bank rating on public sector management and institutions 2018 (CPIA Cluster D)		3.4
Insecticide Resistance Monitoring, Implementation and Impact		
Insecticide classes with mosquito resistance in representative sentinel sites confirmed since 2010	4	
Insecticide resistance monitored since 2015 and data reported to WHO		
National Insecticide Resistance Monitoring and Management Plan		
RDTs in stock (>9 months stock)		
ACTs in stock (>9 months stock)		
LLINIRS campaign on track		
Country Reporting Launch of Zero Malaria Starts with Me Campaign		
Scale of Implementation of ICCM (2017)		
Operational LLINIRS coverage (% of at risk population)		100
On track to reduce case incidence by ≥40% by 2020 (vs 2015)		
On track to reduce case mortality by ≥40% by 2020 (vs 2015)		
Tracer Indicators for Maternal and Child Health and NTDs		
Mass Treatment Coverage for Neglected Tropical Disease (NTD index, %)(2018)		89
Estimated % of Total Population living with HIV who have access to antiretroviral therapy (2019)	▲	67
Estimated % of children (0–14 years old) living with HIV who have access to antiretroviral therapy (2019)	▲	39
% deliveries assisted by skilled birth attendant		80
Postnatal care (within 48 hrs)		74
Exclusive breastfeeding (% children < 6 months)		50
Vitamin A Coverage 2018 (2 doses)		99
DPT3 coverage 2019 (vaccination among 0-11 month olds)		91

Key

	Target achieved or on track
	Progress but more effort required
	Not on track
	No data
	Not applicable

Malaria

Sustaining Essential Health Services During the COVID-19 Pandemic

The COVID-19 pandemic is putting an incredible strain on health systems across Africa. Health systems are required to maintain routine health services for other illnesses even as they handle the additional burden. In order to prevent widespread morbidity and mortality, it is of vital importance that we work to sustain the delivery of essential life-saving interventions during this difficult time including for Reproductive, Maternal, Newborn, Adolescent and Child health and malaria.

WHO underlines the critical importance of sustaining efforts to prevent, detect and treat malaria during the COVID-19 pandemic. It is of vital importance to ensure the continuity of malaria prevention and treatment services including distribution of insecticide-treated nets and indoor residual spraying, as well as chemoprevention for pregnant women and young children (intermittent preventive treatment in pregnancy and seasonal malaria chemoprevention). Any intervention must consider the importance of both lowering malaria-related mortality and ensuring the safety of communities and health workers given the ease of transmission of COVID-19.

For Burkina Faso, it will be of vital importance to ensure that the delivery of seasonal malaria chemoprevention planned for 2020 goes ahead as planned, whilst taking into account physical distancing, in accordance with the recent guidance and recommendations from WHO. Without this campaign, coupled with the need to sustain essential health services including malaria case management, there will be an increase in malaria cases and deaths. Under the worst-case scenario, in which all ITN campaigns are suspended and there is a 75% reduction in access to effective antimalarial medicines, WHO estimate that there could be a 10.3% increase in malaria cases, and a 73.9% increase in malaria deaths in Burkina Faso. This scenario would represent a complete reversal in the substantial progress in malaria mortality reductions seen over the last 2 decades.

Progress

Burkina Faso has scaled up coverage of iCCM and has procured sufficient LLINs to achieve universal operational coverage of the targeted at risk population. Burkina Faso has a high rating in public sector management systems (CPIA cluster D). The country has carried out insecticide resistance monitoring since 2015 and has reported the results to WHO and has recently completed the insecticide resistance monitoring and management plan. Burkina Faso has enhanced the tracking and accountability mechanisms for malaria with the development of the Malaria Scorecard. Burkina Faso has decreased the estimated malaria mortality rate by more than 40% since 2015. The country is also showing leadership in malaria control through participation in the High Burden High Impact approach.

Impact

The annual reported number of malaria cases in 2018 was 11,970,321 with 4,294 deaths.

Key Challenges

- Insecticide resistance threatens vector control effectiveness.
- Sustaining the delivery of essential life-saving interventions during the COVID-19 pandemic including for Reproductive, Maternal, Newborn, Adolescent and Child health including malaria.

Previous Key Recommended Actions

Objective	Action Item	Suggested completion timeframe	Progress	Comments - key activities/accomplishments since last quarterly report
Address funding	Ensure the GF malaria funding application is submitted by Q2 2020 and ensure that resources are allocated to malaria control at a level that is sufficient to sustain the gains made in recent years	Q2 2020		The country submitted the GF malaria application in Q2
Impact	Ensure that malaria services including case management and vector control are sustained and implemented whilst using COVID-19 sensitive guidelines during the pandemic	Q4 2020 Impact		The country has adequate stocks of RDTs and is working to accelerate the delivery of ACTs. The IRS campaign took place as planned, whilst planning is ongoing for the SMC campaigns beginning in Q3 2020

RMNCAH and NTDs

Progress

Burkina Faso has also made good progress in tracer RMNCAH interventions, including DPT3, exclusive breastfeeding and skilled birth attendants. Burkina Faso has significantly enhanced the tracking and accountability mechanisms with the development of the Maternal, Newborn, Child and Adolescent Health Scorecard.

Progress in addressing Neglected Tropical Diseases (NTDs) in Burkina Faso is measured using a composite index calculated from preventive chemotherapy coverage achieved for lymphatic filariasis, onchocerciasis, schistosomiasis and trachoma. Preventive chemotherapy coverage in Burkina Faso is high for schistosomiasis (100%), trachoma (100%), onchocerciasis (82%) and for lymphatic filariasis (77%). Overall, the NTD preventive chemotherapy coverage index for Burkina Faso in 2018 is 89 and shows a slight decrease compared to 2017 index (90).

Previous Key Recommended Actions

Objective	Action Item	Suggested completion timeframe	Progress	Comments - key activities/accomplishments since last quarterly report
RMNCAH ¹ : Impact	Ensure that essential RMNCAH services are sustained and implemented whilst using Covid-19 sensitive guidelines during the pandemic. Address any stock-outs of essential RMNCAH commodities	Q4 2020		Deliverable not yet due

Burkina Faso has responded positively to the RMNCAH recommended action addressing the low coverage of ARTs in children, and continues to track progress as these actions are implemented.

¹ RMNCAH metrics, recommended actions and response tracked through WHO

Key

	Action achieved
	Some progress
	No progress
	Deliverable not yet due